

# Western Australian Immunisation Schedule

## Childhood

Age	Antigen	Vaccine brand(s)	Important information
<b>Birth to &lt;7 days</b>	Hepatitis B	H-B-Vax <sup>®</sup> II Paed or Engerix <sup>®</sup> B Paed	
<b>6 Weeks</b>	DTPa-hepB-IPV-Hib 13vPCV Rotavirus	Infanrix <sup>®</sup> hexa Prevenar <sup>®</sup> 13 Rotarix <sup>®</sup>	Oral dose 1: 6–14 weeks of age
<b>Aboriginal</b>	MenACWY	Nimenrix <sup>®</sup>	
<b>Aboriginal and/or MRC</b>	MenB	Bexsero <sup>®</sup>	<a href="#">Prophylactic paracetamol</a> recommended prior to dose
<b>4 Months</b>	DTPa-hepB-IPV-Hib 13vPCV Rotavirus	Infanrix <sup>®</sup> hexa Prevenar <sup>®</sup> 13 Rotarix <sup>®</sup>	Oral dose 2: 10–24 weeks of age
<b>Aboriginal</b>	MenACWY	Nimenrix <sup>®</sup>	
<b>Aboriginal and/or MRC</b>	MenB	Bexsero <sup>®</sup>	<a href="#">Prophylactic paracetamol</a> recommended prior to dose
<b>6 Months</b>	DTPa-hepB-IPV-Hib	Infanrix <sup>®</sup> hexa	
<b>Aboriginal and/or MRC</b>	13vPCV	Prevenar <sup>®</sup> 13	
<b>MRC</b>	MenB	Bexsero <sup>®</sup>	<a href="#">Prophylactic paracetamol</a> recommended prior to dose
<b>12 Months</b>	MMR 13vPCV MenACWY	MMRII <sup>®</sup> or Priorix <sup>®</sup> Prevenar <sup>®</sup> 13 Nimenrix <sup>®</sup> *	
<b>Aboriginal and/or MRC</b>	MenB	Bexsero <sup>®</sup>	<a href="#">Prophylactic paracetamol</a> recommended prior to dose
<b>MRC</b>	Hepatitis B	H-B-Vax <sup>®</sup> II Paed or Engerix <sup>®</sup> B Paed	<32 weeks gestation or <2000g birth weight
<b>&gt;12 Months</b>	At risk children >12 months of age require 1 dose of 13vPCV at diagnosis followed by 2 doses of 23vPPV. See the <a href="#">Australian Immunisation Handbook</a> for more information.		
<b>MRC</b>			
<b>18 Months</b>	DTPa Hib MMRV	Infanrix <sup>®</sup> or Tripacel <sup>®</sup> ActHIB <sup>®</sup> Priorix-Tetra <sup>®</sup> or ProQuad <sup>®</sup>	Do not administer MMRV as dose 1 of MMR containing vaccine if <4 years of age
<b>Aboriginal</b>	Hepatitis A	Vaqa <sup>®</sup> Paed/Adol	First dose of the 2 dose schedule
<b>&lt;2 Years</b>	Catch-up for MenB is available for Aboriginal children <2 years of age until 30 June 2023.		
<b>Aboriginal</b>			
<b>4 Years</b>	DTPa-IPV	Quadracel <sup>®</sup> or Infanrix <sup>®</sup> IPV	
<b>Aboriginal</b>	Hepatitis A	Vaqa <sup>®</sup> Paed/Adol	Not required if previously received 2 doses (first dose at age ≥12 months) at least 6 months apart
<b>Aboriginal and/or MRC</b>	23vPPV	Pneumovax23 <sup>®</sup>	Subsequent dose of 23vPPV at least 5 years later

## Adolescent and adult

Age	Antigen	Vaccine brand(s)	Important information
<b>All adolescents and adults with MRC</b>	At risk persons >12 months of age require 1 dose of 13vPCV at diagnosis followed by 2 doses of 23vPPV. People of all ages with specified MRC are eligible for MenACWY, MenB and Hib vaccines. See the ATAGI clinical advice for <a href="#">pneumococcal</a> and <a href="#">meningococcal</a> diseases as well as the <a href="#">Australian Immunisation Handbook</a> for more information.		
<b>Year 7</b>	HPV dTpa	Gardasil®9 Boostrix®	2 doses of Gardasil®9 6 months apart
<b>Year 10</b>	MenACWY	Nimenrix®	Catch-up for adolescents <20 years of age
<b>Pregnant</b>	dTpa	Boostrix® or Adacel®	Each pregnancy, ideally between 20–32 weeks
<b>≥50 Years Aboriginal</b>	13vPCV + 23vPPV	Prevenar® 13 + Pneumovax23®	See the <a href="#">Australian Immunisation Handbook</a> for for timing of doses
<b>Born after 1965</b>	MMR	MMRII® or Priorix®	Give 1 or 2 doses as required
<b>≥70 Years</b>	13vPCV	Prevenar® 13	
<b>≥70 to 79 Years</b>	Herpes-zoster	Zostavax®	Catch-up program for 71–79 year olds until 31 Oct 2021

## 2020 Seasonal influenza

Age	Vaccine brand(s)	Important information
<b>6 Months to &lt;5 Years</b>	Vaxigrip® Tetra, Fluarix® Tetra, FluQuadri®	If child <9 years, 2 doses required 1 month apart in the first year they receive influenza vaccine
<b>≥5 Years to Year 6</b>	FluQuadri® (WA State Program)	
<b>5 to &lt;65 Years Aboriginal and/or MRC</b>	Afluria® Quad, Fluarix® Tetra, FluQuadri®, Vaxigrip® Tetra	
<b>Pregnant</b>	Afluria® Quad, Fluarix® Tetra, FluQuadri®, Vaxigrip® Tetra	Each pregnancy, any trimester
<b>≥65 Years</b>	Fluad® Quad	Use standard QIV if Fluad® Quad unavailable
<b>WA Health staff</b>	Fluarix® Tetra, FluQuadri® (WA State Program)	For ≥65 years use Fluad® Quad

The term 'Aboriginal' is inclusive of all Aboriginal and Torres Strait Islander people.

MRC refers to medical risk conditions. Refer to the current online edition of The [Australian Immunisation Handbook](#) for all medical risk conditions.

Persons aged <20 years can catch-up scheduled vaccines, refer to [Australian Immunisation Handbook](#) for timing of doses.

Adult refugees and humanitarian entrants are eligible for dTpa, Hepatitis B, IPV, MMR and VZV.

\* MenACWY catch-up for Aboriginal children <5 years of age.